

CSS Past Paper Applied Mathematics (2018)

For a comprehensive collection of CSS preparation resources; date sheets, notes, solved past papers, examiner reports, and FPSC-recommended Books, please visit our website or feel free to reach out to us. We are here to assist you in your CSS journey.





FEDERAL PUBLIC SERVICE COMMISSION COMPETITIVE EXAMINATION-2018 FOR RECRUITMENT TO POSTS IN BS-17 UNDER THE FEDERAL GOVERNMENT

APPLIED MATHEMATICS

| TIME ALLOWED: THREE HOURS | MAXIMUM MARKS = 100 |
|-----------------------------------|---------------------|
| · · · · · · · · · · · · · · · · · | |

- NOTE:(i) Attempt ONLY FIVE questions. ALL questions carry EQUAL marks(ii) All the parts (if any) of each Question must be attempted at one place instead of at different places.
 - (iii) Candidate must write Q. No. in the Answer Book in accordance with Q. No. in the Q.Paper.
 - (iv) No Page/Space be left blank between the answers. All the blank pages of Answer Book must be crossed.
 - (v) Extra attempt of any question or any part of the attempted question will not be considered.
 - (vi) Use of Calculator is allowed.

Q. No. 1. (a) If
$$\psi = Sin \frac{(kr)}{r}$$
, then show that $\nabla^2 \psi + k^2 \psi = 0$. (10)

(b) Calculate the Line Integral $\int_c A.dr$, where $A = \frac{-yi + xj}{x^2 + y^2}$, and the curve C is (10)

given by the equations $x^2 + y^2 = a^2$ and Z = 0.

- Q. No. 2. (a) Forces of magnitude P, 2P, 3P, 4P act respectively along the sides AB, BC, CD, (10) DA of a square ABCD, of sides a, and forces each of magnitude (8√2) P act along the diagonals BD, AC. Find the magnitude of the resultant force and distance of its line of action from A.
 - (b) A uniform ladder, of length 70 feet, rests against a vertical wall with which it makes an angle of 45° , the coefficient of friction between the ladder and the wall and the ground respectively being $\frac{1}{3}$ and $\frac{1}{2}$. If a man, whose weight is one half that of the ladder, ascends the ladder, where will he be when the ladder slips?
- Q. No. 3. (a) A particle moves in a straight line with an acceleration kv^3 . If its initial velocity is (10) u, find the velocity and the time spent when the particle has travelled a distance x.
 - (b) Derive the Tangential and Normal components of the velocity and acceleration. (10)
- Q. No. 4. (a) Solve the following Cauchy- Euler Equation

$$x^2 \frac{d^2 y}{dx^2} - 2x \frac{dy}{dx} - 4y = 0.$$

(b) Convert the following Bernoulli Differential Equation into standard form and (10) then solve.

$$\frac{dy}{dx} + \left(\frac{xy}{1-x^2}\right) = xy^{\frac{1}{2}}.$$

Q. No. 5. (a) Convert the following Ordinary Differential Equation into standard form and then (10) solve using Method of Variation of Parameters.

$$x^2y'' - 3xy' + 3y = 2x^4e^x$$

(b) Check whether the following Ordinary Differential Equation is an Exact Equation (10) or not. If yes, then solve.

$$(3x^2y + 2)dx + (x^3 + y)dy = 0$$

Page 1 of 2

(10)

APPLIED MATHEMATICS

Q. No. 6. (a) Find the Fourier Series of f on the given interval.

$$f(x) = \begin{cases} -1, & -\pi < x < 0\\ 1, & 0 \le x < \pi \end{cases}$$

(b) Solve the following Partial Differential Equation subject to the conditions given. (10)

$$a^2 \frac{\partial^2 u}{\partial x^2} = \frac{\partial^2 u}{\partial t^2}, \qquad 0 < x < L, \qquad t > 0,$$

$$u(0,t) = 0,$$
 $u(L,t) = 0,$ $t > 0,$

$$u(x,0) = f(x), \ \frac{\partial u}{\partial t} = g(x)[at \ time \ t = 0], \qquad and \quad 0 < x < L.$$

Q. No. 7. (a) Use Newton-Raphson method to find solution accurate to within 10^{-4} for the nonlinear equation. (10)

$$x^3 - 2x^2 - 5 = 0, I = [1,4]$$

(b) Use Lagrange Interpolating polynomial of degree two to approximate f(8.4), If (10)

$$f(8.1) = 16.94410$$
, $f(8.3) = 17.56492$, $f(8.6) = 18.50515$, $f(8.7) = 18.82091$

Q. No. 8. (a) Approximate $\int_0^1 \frac{dx}{1+x^2}$ using Trapezoidal rule and Simpson's rule with n=4. (10) Also compare your results with the exact value of the integral.

(b) Use Euler's method to approximate the solution of the following initial value (10) problem.

$$y' = \frac{1+y}{t}$$
, $1 \le t \le 2$, with $y(1) = 2$, $h = 0.25$

(10)

Reach out to us @ <u>info@thinked.co</u> If you are interested in writing for us email us at <u>writeforthinked@thinked.co</u>